

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

DS

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : A01N 51/00 // (A01N 51/00, 43:828)		A1	(11) International Publication Number: WO 00/05959
		(43) International Publication Date:	10 February 2000 (10.02.00)
(21) International Application Number: PCT/EP99/05414		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 28 July 1999 (28.07.99)			
(30) Priority Data: 9816638.2 30 July 1998 (30.07.98) GB			
(71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).			
(71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).			
(72) Inventors; and (75) Inventors/Applicants (for US only): OOSTENDORP, Michael [DE/DE]; Uhlendweg 7, D-79618 Rheinfelden (DE). RUESS, Wilhelm [CH/CH]; Burgweg 4, CH-4148 Pfeffingen (CH).			
(74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).			
		Published With international search report Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: PESTICIDAL COMPOSITIONS COMPRISING 1,2,3-BENZOTHIADIAZOLE DERIVATIVES			
<p>(1)</p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p> <p>(f)</p> <p>(g)</p> <p>(h)</p> <p>(i)</p>			
(57) Abstract			
<p>A composition and method for protecting and immunizing plants against attack by insects, acarina and nematodes, comprising at least two active ingredient components in amounts having synergistic action, together with a suitable carrier, wherein component I is a compound of formula (I) in which Het is an optionally substituted 5- or 6-membered aromatic ring having 1-3 hetero atoms N, O and/or S; in particular selected from the group (a, b, c, d, e, f, g, h, i) wherein R₁ is hydrogen, optionally substituted C₁-C₆alkyl, optionally substituted C₂-C₆alkenyl or optionally substituted phenyl and R₂ and R₃ independently of each other are hydrogen or optionally substituted C₁-C₆alkyl, and acid addition salts and metal salt complexes thereof; and wherein component II is selected from the insecticides IIA) imidacloprid and IIB) thiamethoxam.</p>			

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing International applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroun	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PESTICIDAL COMPOSITIONS COMPRISING 1,2,3-BENZOTHIADIAZOLE DERIVATIVES


The present invention relates to new compositions and methods for protecting and immunizing plants against attack by insects, acarina and nematodes.

- 5 The compositions comprise at least two active ingredient components in an effective amount together with a suitable carrier, wherein component I is a compound of formula I



in which

Het is an optionally substituted 5- or 6-membered aromatic ring having 1-3 hetero atoms N, O and/or S; in particular selected from the group



wherein

R₁ is hydrogen, optionally substituted C₁-C₆alkyl, optionally substituted C₂-C₆alkenyl or optionally substituted phenyl and

R₂ and R₃ independently of each other are hydrogen or optionally substituted C₁-C₆alkyl, and acid addition salts and metal salt complexes thereof (EP-A-816,363);

and wherein component II is selected from the insecticides

IIA) imidacloprid (The Pesticide Manual, 11th. Ed. 1997, No.418)

and

IIb) thiamethoxam (5- (2-chlorothiazol-5-ylmethyl)-3-methyl-4-nitroimino-perhydro-1,3,5-oxadiazine) (EP-A-580,553).

It is known from EP-A-816,363 that compounds of formula I are active as microbicides and may be used in agriculture and horticulture for directly combating undesired phytopathogenic fungi and bacteria and for immunizing plants against attack of phytopathogenic fungi and bacteria by inducing "Systemic Activated Resistance" (SAR) in the plants. Surprisingly it has now been found that compositions and methods according to the invention exhibit synergistic activities and are therefore particularly effective for protecting and immunizing plants against attack by insects, acarina and nematodes.

In the above formula alkyl designates straight chain and branched alkyl groups, such as methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, i-butyl, t-butyl, n-pentyl, i-pentyl, s-pentyl, neo-pentyl, and the various isomers of hexyl.

Alkenyl is straight-chain or branched alkenyl such as allyl, methallyl, 1-methylvinyl or but-2-en-1-yl. Preferred alkenyl radicals contain 3 to 4 carbon atoms in the chain.

Substituents at alkyl, alkenyl and phenyl include halogen, such as fluoro, chloro, bromo and iodo; cyano; alkoxy, such as methoxy, ethoxy, n-propoxy, i-propoxy, and the various isomers of butoxy; alkylthio, such as methylthio and ethylthio; alkoxy-carbonyl, such as methoxycarbonyl and ethoxycarbonyl; and phenyl.

Preferred compounds of formula I are those wherein

R₁ is hydrogen, C₁-C₄alkyl, C₁-C₄haloalkyl, phenyl-C₁-C₄alkyl, cyano- C₁-C₄alkyl, C₁-C₂alkoxy-C₁-C₄alkyl, C₁-C₂alkylthio- C₁-C₄alkyl, C₁-C₂alkoxycarbonyl- C₁-C₄alkyl, C₂-C₄alkenyl, phenyl, halogen substituted phenyl or C₁-C₄alkyl substituted phenyl and R₂ and R₃ independently of each other are hydrogen or C₁-C₄alkyl.

Particularly preferred are those, wherein

R₁ is hydrogen, methyl, ethyl, propyl, isopropyl, butyl, cyanomethyl, 2-cyanoethyl, 1-cyanoethyl, 3-cyano-propyl, methoxymethyl, ethoxymethyl, methylthiomethyl, methoxycarbonyl-ethyl, ethoxycarbonyl-methyl, ethoxycarbonyl-ethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, benzyl, phenyl, chlorophenyl, methylphenyl, vinyl, allyl or but-3-en-1-yl, and

R₂ and R₃ independently of each other are hydrogen, methyl, ethyl, propyl or isopropyl.

Especially preferred are the specific compounds as disclosed in EP-A-816,363.

Suitable acid addition salts and metal salts complexes thereof are also described in this reference.

The compositions of the invention can be used in the agricultural sector and related fields preventively and/or curatively. Plants can be protected both by direct action on the pests or by activation and stimulation of the plant's own defense system (immunization). The latter mode of action has also become known by the name "Systemic Activated Resistance" ("SAR").

Accordingly, using the mixtures of the invention, it is possible to control plant diseases on the one hand by strengthening the plant by activating its own defence system and on the other hand by additionally controlling the pathogens directly.

A particular advantage of the mixtures according to the invention is further that, because the modes of action of components I and II are completely different, the threat of resistance being developed in the control of plant diseases is effectively prevented.

The compositions can also be used as dressings in the treatment of seed (fruit, tubers, grains) and plant cuttings to provide protection.

Advantageous mixing ratios of the two active ingredients are

I:II = 200:1 to 1:100, 50:1 to 1:20, 10:1 to 1:5, and preferably 10:1 to 1:1.

The compositions are effective against the following phytopathogenic pests:

A) Insects, for example Lepidoptera (e.g. *Cydia*, *Heliothis*, *Lobesia*, *Pyralidae*, *Geometridae*, *Noctuidae*); Thysanoptera (e.g. *Frankliniella*, *Thrips palmi*); Heteroptera (e.g. *Capsidae*, *Pentatomidae*); Homoptera (e.g. *Bemisia tabaci*, *Nilaparvata*, *Delphacidae*, *Jassidae*, *Psyllina*, *Aphidina*, *Aphis*, *Coccina*); Diptera (e.g. *Musca*, *Trypetidae*, *Cecidomyiidae*); Coleoptera (e.g. *Chrysomelidae*, *Curculionidae*, *Lissorhoptus*); Cicadellidae (e.g. *Nephotettix*); Cryptophagidae (e.g. *Atomaria*); Erisiornathidae (e.g. *Pemphigus*).

B) Acarina, for example *Tetranychidae*, *Tarsonemidae*, *Eriophyidae*, *Phyllocoptidae*;

C) Nematodes, for example of the genera *Heterodera* and *Globodera* (cystogenic nematodes), *Meloidogyne* (root-knot nematodes), *Radopholus*, *Pratylenchus*, *Tylenchulus*, *Longidorus*, *Trichodorus*, *Xiphinema*, *Ditylenchus* (stem parasites), *Aphelenchoides* (leaf

nematodes), and Anguina (blossom nematodes); particularly harmful nematode species of the genus Meloidogyne, for example Meloidogyne incognita, and of the genus Heterodera, for example Heterodera glycines (soybean cyst nematode).

Target crops to be activated within the scope of the present invention comprise e.g. the following species of plants: cereals (wheat, barley, rye, oats, rice, maize, sorghum and related species); beet (sugar beet and fodder beet); pomes, stone fruit and soft fruit (apples, pears, plums, peaches, almonds, cherries, mirabella, mangos strawberries, raspberries and blackberries); leguminous plants (beans, lentils, peas, soybeans); oil plants (rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans, groundnuts); cucumber plants (marrows, cucumber, melons); fibre plants (cotton, flax, hemp, jute); citrus fruit (oranges, lemons, grapefruit, mandarins); vegetables (spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, paprika); lauraceae (avocados, cinnamon, camphor); and plants such as tobacco, nuts, coffee, aubergines, sugar cane, tea, pepper, vines, hops, bananas and natural rubber plants, as well as ornamentals (flowers, shrubs, broad-leaved trees and evergreens, such as conifers). This list does not represent any limitation.

The compositions are particularly useful for treating

- rice, in particular for seedling box-, into water granules- and for foliar application;
- potatoes and tomatoes, in particular for foliar application;
- sugar beet, in particular for seed treatment.

The method of the invention comprises applying to the plants to be treated or the locus thereof in admixture or separately, an effective amount of a component I and a component II in any desired sequence or simultaneously.

The term locus as used herein is intended to embrace the fields on which the treated crop plants are growing, or where the seeds of cultivated plants are sown, or the place where the seed will be placed into the soil. The term seed is intended to embrace plant propagating material such as cuttings, seedlings, seeds, germinated or soaked seeds.

A method of applying the composition is application to the leaves (foliar application). The frequency and rate of application depend upon the risk of infestation by the corresponding pathogen. The compounds can also penetrate the plant through the roots via the soil

(systemic action) if the locus of the plant is impregnated with a liquid formulation or if the substances are introduced in solid form into the soil, e.g. in the form of granules (soil application). In paddy rice crops, such granules can be applied in metered amounts to the flooded rice field or to the seedling box before transplanting. In order to treat seed, the compounds I can, however, also be applied to the seeds (coating), either by impregnating the grains or tubers with a liquid formulation of the active ingredient, or by coating them with a solid formulation.

The compounds are used together with the adjuvants conventionally employed in formulation technology. For that purpose they are advantageously formulated in known manner e.g. into emulsifiable concentrates, coatable pastes, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granules or by encapsulation in e.g. polymer substances. As with the nature of the compositions, the methods of application, such as spraying, atomizing, dusting, scattering, coating or pouring, are chosen in accordance with the intended objectives and the prevailing circumstances. Advantageous rates of application of the active ingredient mixture are normally from 0.01 to 10 kg of active ingredient (a.i.) per hectare, preferably from 30 g to 1000 g a.i./ha, especially from 50 g to 500 g a.i./ha. In the case of the treatment of seed, the rates of application are from 0.5 g to 1000 g, preferably from 5 g to 100 g, a.i. per 100 kg of seed. The agrochemical compositions generally comprise 0.1 to 99 % by weight, preferably 0.1 to 95 % by weight, of a compound of formula I, 99.9 to 1 % by weight, preferably 99.8 to 5 % by weight, of a solid or liquid adjuvant and 0 to 25 % by weight, preferably 0.1 to 25 % by weight, of a surfactant. The compositions may also comprise further auxiliaries, such as stabilizers, antifoams, viscosity regulators, binders or tackifiers, as well as fertilizers or other active ingredients for obtaining special effects. Manufacturing of the formulations is a routine matter in the art of pesticides and largely depends on the desired mode of application and target plant.

ExamplesFormulation ExamplesWettable powders

	a)	b)	c)
active ingredient (I:II = 3:1)	25 %	50 %	75 %
sodium lignosulfonate	5 %	5 %	-
sodium lauryl sulfate	3 %	-	5 %
sodium diisobutyl naphthalenesulfonate	-	6 %	10 %
phenol polyethylene glycol ether (7-8 mol of ethylene oxide)	-	2 %	-
highly dispersed silicic acid	5 %	10 %	10 %
kaolin	62 %	27 %	-

The active ingredient is thoroughly mixed with the adjuvants and the mixture is thoroughly ground in a suitable mill, affording wettable powders which can be diluted with water to give suspensions of the desired concentration.

Emulsifiable concentrate

active ingredient (I : II) = 4:1	10 %
octylphenol polyethylene glycol ether (4-5 mol of ethylene oxide)	3 %
calcium dodecylbenzenesulfonate	3 %
castor oil polyglycol ether (35 mol of ethylene oxide)	4 %
cyclohexanone	30 %
xylene mixture	50 %

Emulsions of any required dilution, which can be used in plant protection, can be obtained from this concentrate by dilution with water.

Extruder granules

active ingredient (I : II = 2:1)	15 %
sodium lignosulfonate	2 %
carboxymethylcellulose	1 %
kaolin	82 %

The active ingredient is mixed and ground with the adjuvants, and the mixture is moistened with water. The mixture is extruded and then dried in a stream of air.

Coated granules

active ingredient (I II = 2:1)	8 %
polyethylene glycol (mol. wt. 200)	3 %
kaolin	89 %

The finely ground active ingredient is uniformly applied, in a mixer, to the kaolin moistened with polyethylene glycol. Non-dusty coated granules are obtained in this manner.

Slow Release Capsule Suspension

28 parts of a combination of the compounds I and II or of each of these compounds separately, are mixed with 2 parts of an aromatic solvent and 7 parts of toluene dilsocyanate/polymethylene-polyphenylisocyanate-mixture (8:1). This mixture is emulsified in a mixture of

1.2 parts of polyvinylalcohol, 0.05 parts of a defoamer and 51.6 parts of water until the desired particle size is achieved. To this emulsion a mixture of 2.8 parts 1,6-diaminohexane in 5.3 parts of water is added. The mixture is agitated until the polymerisation reaction is completed.

The obtained capsule suspension is stabilized by adding 0.25 parts of a thickener and 3 parts of a dispersing agent. The capsule suspension formulation contains 28% of the active ingredients. The medium capsule diameter is 8-15 microns.

Seed Dressing Formulation

25 parts of a combination of the compounds I and II, 15 parts of dialkylphenoxypoly(ethylenoxy)ethanol, 15 parts of fine silica, 44 parts of fine kaolin, 0.5 parts of Rhodamine B as a colorant and 0.5 parts of Xantham Gum are mixed and ground in a contraplex mill at approx. 10000 rpm to an average particle size of below 20 microns. The resulting formulation is applied to seeds as an aqueous suspension in an apparatus suitable for that purpose.

Biological examples:

A synergistic effect exists whenever the action of an active ingredient combination is greater than the sum of the actions of the individual components.

The action to be expected E for a given active ingredient combination obeys the so-called COLBY formula and can be calculated as follows:

ppm = milligrams of active ingredient (= a.i.) per litre of spray mixture

X = % action by active ingredient I using p ppm of active ingredient

Y = % action by active ingredient II using q ppm of active ingredient.

According to Colby, the expected (additive) action of active ingredients I+II using p+q ppm

of active ingredient is $E = X + Y - \frac{X \cdot Y}{100}$

If the action actually observed (O) is greater than the expected action (E), then the action of the combination is superadditive, i.e. there is a synergistic effect.

O/E = synergy factor (SF).

In the following examples the compositions exhibit synergistic action:

Example B-1.1: Action against *Aphis craccivora* (insect)

Pea seedlings are treated with a spray mixture comprising 100 ppm of a mixture of active ingredients and 3 days later infected with *Aphis craccivora* and incubated at 20-22°C. 6 and 12 days later, the percentage reduction in population (% activity) is determined.

Example B-2: Action against *Liriomyza* sp. (insect)

After a cultivation period of 3 weeks, tomato plants are sprayed with a spray mixture prepared from a wettable powder formulation of the test compounds. After 96 hours, the treated plants transferred into large buckets sealed with cellophane. Adult *Liriomyza* are introduced into each bucket and allowed to feed and oviposit for 24 h and then removed. After 13-17 days the experiments are evaluated by counting number of mines, oviposition and feeding punctures.

Example B-3: Action against *Bemisia argentifolii* (insect)

After a cultivation period of 3 weeks, tomato plants are sprayed with a spray mixture prepared from a wettable powder formulation of the test compounds. After 96 hours, the treated plants covered under closed cylinders and *Bemisia* is introduced. Eggs and nymphs are counted on the terminal trifoliate after 5 to 10 days using a stereomicroscope. In addition plant damage is estimated visually.

Example B-4: Action against Tetranychus urticae (Acarina)

Young bean plants are sprayed with an aqueous emulsion spray mixture which comprises 400 ppm of the active ingredients; one day later the plants are populated with a mixed population of Tetranychus urticae and subsequently incubated for 6 days at 25°C. The percentage reduction in population (% action) is determined by comparing the number of dead eggs, larvae and adults on the treated with those on the untreated plants.


Example B-5: Action against a mixed population of Meloidogyne javanica and Meloidogyne arenaria (Nematodes) on grapevines.

Grapevines (Cabernet Sauvignon) are treated with a spray mixture prepared from a wettable powder formulation of the test compounds one week before inoculation, at inoculation and/or one week after inoculation with a mixed population of Meloidogyne javanica and Meloidogyne arenaria.

After 4 weeks, the total number of eggs produced is evaluated.


Claims

1. A composition for protecting and immunizing plants against attack by insects, acarina and nematodes, comprising at least two active ingredient components in amounts having synergistic action, together with a suitable carrier, wherein component I is a compound of formula



In which

Het is an optionally substituted 5- or 6-membered aromatic ring having 1-3 hetero atoms N, O and/or S; in particular selected from the group



wherein

R₁ is hydrogen, optionally substituted C₁-C₆alkyl, optionally substituted C₂-C₆alkenyl or optionally substituted phenyl and

R₂ and R₃ independently of each other are hydrogen or optionally substituted C₁-C₆alkyl, and acid addition salts and metal salt complexes thereof; and wherein component II is selected from the insecticides

IIA) imidacloprid and IIB) thiamethoxam.

2. A method of protecting plants against attack by insects, acarina and nematodes by treating the plants, parts of the plants or their locus with an effective amount of component I and component II according to claim 1, in any desired sequence or simultaneously.

3. A method according to claim 2 wherein rice, potatoes, tomatoes and sugar beet are treated.

4. A method according to claim 2 wherein seed is treated.
